

Cambridge International A Level

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

May/June 2024

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **20** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.


Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1(a)		B1	<p>Correct shape, roughly symmetrical. Both sections should be solid straight lines. If not drawn with a ruler the intention must be clear. Allow construction lines if dashed or clearly fainter. $2a$ marked on each axis (must be $2a$, not just 2). Needs to extend into negative x. If a is given a value, then B0. Ignore $y = 2x - 3a$ if seen.</p>
		1	
1(b)	Solve linear equation or inequality to obtain critical value $x = \frac{5}{3}a$ or exact equivalent.	B1	Ignore $x = a$ if seen.
	Obtain $x < \frac{5}{3}a$ or exact equivalent	B1	<p>Accept $x < \frac{10}{6}a$ or $(-\infty, \frac{5}{3}a)$. Must be strict inequality. Need a clear final solution: $x > a$ or $x < a$ must be rejected if seen as part of the working. Rejection can be implied, e.g. if only the correct inequality is underlined. B0 B0 if a is given a value.</p>
Alternative Method for Question 1(b)			
	Solve quadratic equation $(2x - 3a)^2 = (x - 2a)^2$ to obtain critical value $x = \frac{5}{3}a$ or exact equivalent	(B1)	<p>$(3x^2 - 8ax + 5a^2 = 0)$ Ignore $x = a$ if seen.</p>
	Obtain $x < \frac{5}{3}a$ or exact equivalent	(B1)	<p>Accept $x < \frac{10}{6}a$ or $(-\infty, \frac{5}{3}a)$. Must be strict inequality. Need a clear final solution: $x > a$ or $x < a$ must be rejected if seen as part of the working. Rejection can be implied, e.g. if only the correct inequality is underlined. B0 B0 if a is given a value.</p>
		2	

PUBLISHED

Question	Answer	Marks	Guidance
2	State or imply the form $A + \frac{B}{2x+3} + \frac{C}{x-4}$	B1	$\frac{Dx+E}{2x+3} + \frac{F}{x-4}$ and $\frac{P}{2x+3} + \frac{Qx+R}{x-4}$ are also valid.
	Use a correct method for finding a constant	M1	SC: If score B0, they can score M1 A1 for one correct constant. B0 M1 A0 available if they substitute two values to form simultaneous equations but get an incorrect answer, or they substitute one value and make an arithmetic error.
	Obtain one of $A = 3$, $B = -2$ and $C = 4$	A1	SC: If the horizontal equation is correct apart from an incorrect value for A , the other A marks may be available.
	Obtain a second value	A1	SC: If denominator factorised as $(x + \frac{3}{2})(x - 4)$ can score a maximum of B0 M1 A1 A1 A0 for a split involving 3 terms.
	Obtain a third value	A1	ISW Statement of the final split is not required.

PUBLISHED

Question	Answer	Marks	Guidance
2	Alternative method for Question 2		
	Divide numerator by denominator	(M1)	
	Obtain $3 \left(+ \frac{Px+Q}{2x^2-5x-12} \right)$	(A1)	$\left(3 + \frac{6x+20}{(2x+3)(x-4)} \right)$
	State or imply the form $\frac{Px+Q}{2x^2-5x-12} = \frac{D}{2x+3} + \frac{E}{x-4}$	(B1)	Must deal with the 3 separately or include it correctly on both sides in their split.
	Obtain one of $D = -2$ and $E = 4$	(A1)	SC: If denominator factorised as $\left(x + \frac{3}{2}\right)(x-4)$, then can score a maximum of B0 M1 A1 A1 A0 for a split involving three terms.
	Obtain a second value	(A1)	ISW Statement of the final split is not required.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
3(a)	Use logarithms to obtain a correct expression without powers e.g. $(2y - 1)\ln a = (x - y)\ln b$	B1	Could use logs to any base e.g. $2y - 1 = (x - y)\log_a b$. Do not condone missing brackets unless recovered later.
	Separate terms and factorise to obtain $y(2\ln a + \ln b) = x\ln b + \ln a$	B1	Or equivalent, e.g. $y = x\frac{\ln b}{\ln a^2 b} + \frac{\ln a}{\ln a^2 b}$ or $y(2 + \log_a b) = x\log_a b + 1$.
	Clear explanation of linear form. From correct work only.	B1	E.g. equation matches the linear form $y = mx + c$ or $py = qx + r$. Condone if they compare with $y = mx + c$, but do not actually state that it must therefore be a straight line. Stating “this is a linear equation” without comparing to a relevant standard form scores B0. B0 if they have $m = \dots$ and $c = \dots$ correct but never actually mention $y = mx + c$.
		3	
3(b)	Use $a = b^3$ and log laws to simplify their equation	M1	$\left[y = x\frac{\ln b}{\ln b^7} + \frac{\ln b^3}{\ln b^7} \right]$ Denominator reduced to a single log term.
	Obtain $y = \frac{1}{7}x + \frac{3}{7}$	A1	Accept $y = \frac{x}{7} + \frac{3}{7}$ but not $y = \frac{x+3}{7}$.
	Alternative method for Question 3(b)		
	Use $a = b^3$ to obtain $b^{3(2y-1)} = b^{x-y}$ or equivalent	(M1)	Or $\log_a b = \frac{1}{3}$.
	Obtain $y = \frac{1}{7}x + \frac{3}{7}$	(A1)	Accept $y = \frac{x}{7} + \frac{3}{7}$ but not $y = \frac{x+3}{7}$.
	2		

PUBLISHED

Question	Answer	Marks	Guidance
4	Substitute $y = 1$ and obtain $e^x = a$, where $a > 0$	M1	Must come from a quadratic in e^x . $(e^{2x} + e^x - 6 = 0)$ Ignore any negative solution.
	Obtain $e^x = 2$ only	A1	Or equivalent e.g. $x = \ln 2$. Condone $x = 0.693\dots$
	State or imply $\frac{d}{dx}(ye^{2x}) = 2ye^{2x} + e^{2x} \frac{dy}{dx}$	B1	Accept y' for $\frac{dy}{dx}$.
	State or imply $\frac{d}{dx}(y^2e^x) = y^2e^x + 2ye^x \frac{dy}{dx}$	B1	Accept y' for $\frac{dy}{dx}$.
	Differentiate RHS of given equation to obtain zero (could be implied by subsequent work), substitute for x and y and obtain $\frac{dy}{dx} = \dots$	M1	Independent. $\left[2 \times 4 + 4 \frac{dy}{dx} + 2 + 4 \frac{dy}{dx} = 0 \right]$ NB: Could also rearrange to obtain $\frac{dy}{dx}$ then substitute. Both steps are needed for M1.
	Obtain $-\frac{5}{4}$ or -1.25	A1	Correct answer from correct working only Accept $-\frac{10}{8}$. -1.2499 is A0.

PUBLISHED

Question	Answer	Marks	Guidance
4	Alternative method for Question 4: Dividing through by e^x		
	Substitute $y = 1$ and obtain $e^x = a$, where $a > 0$	(M1)	Must come from a quadratic in e^x . $(e^{2x} + e^x - 6 = 0)$ Ignore any negative solution.
	Obtain $e^x = 2$ only	(A1)	Or equivalent e.g. $x = \ln 2$. Condone $x = 0.693\dots$
	State or imply $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$	(B1)	Accept y' for $\frac{dy}{dx}$.
	State or imply $\frac{d}{dx}(ye^x) = ye^x + e^x \frac{dy}{dx}$	(B1)	Accept y' for $\frac{dy}{dx}$.
	Differentiate RHS of given equation to obtain $-6e^{-x}$, substitute for x and y and obtain $\frac{dy}{dx} = \dots$	(M1)	Independent. $\left[1 \times 2 + 2 \frac{dy}{dx} + 2 \frac{dy}{dx} = -\frac{6}{2} \right]$ NB: Could also rearrange to obtain $\frac{dy}{dx}$ then substitute. Both steps are needed for M1.
	Obtain $-\frac{5}{4}$ or -1.25	(A1)	Correct answer from correct working only Accept $-\frac{10}{8}$. -1.2499 is A0.
		6	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	Calculate the value of a relevant expression or values of a pair of expressions at $x=0.7$ and $x=0.8$	M1	Allow if working with a smaller interval, e.g. (0.72, 0.78). Need all relevant values but condone one error. Pairings must be clear for solutions involving four values (do not accept embedded values). M0 if working in degrees e.g -1.94..., 1.04...
	Complete the argument correctly with correct calculated values. (can be using the equation in the rubric or the equation in (b) or equivalent)	A1	E.g. $4.95... > 4.26...$ and $4.06... < 4.49...$ $-0.439... < 0$, $0.690... > 0$ $1.4 < 1.5029...$, $1.6 > 1.449...$ $1.1 > 1$, $0.86 < 1$ $0.0515 > 0$, $-0.075 < 0$. Allow values rounded or truncated to 2sf.
		2	
5(b)	State $2x = \ln(5 + \cos 3x)$ and take exponential of both sides to obtain $e^{2x} = 5 + \cos 3x$	B1	Given answer requires fully correct working or work vice versa. If working in reverse, must get to the iterative formula, including subscripts.
		1	
5(c)	Use the iterative process correctly at least once	M1	M0 if working in degrees (e.g. values heading for 0.89...).
	Obtain final answer 0.740	A1	
	Show sufficient iterations to at least 5dp to justify 0.740 to 3dp, or show that there is a sign change in the interval (0.7395, 0.7405)	A1	E.g. 0.7, 0.75150, 0.73719, 0.74105, 0.74000, 0.74028 0.75, 0.73759, 0.74094, 0.74003, 0.74028 0.8, 0.72494, 0.74443, 0.73909, 0.74053, 0.74014, 0.74025 Allow recovery. Allow truncation or rounding and condone small differences in the final decimal place.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
6(a)	Use correct product rule	*M1	Or equivalent. Condone incorrect chain rule. M0 if a value is used for a (not equivalent work).
	Obtain correct derivative	A1	E.g. $\frac{dy}{dx} = -axe^{-ax} + e^{-ax}$
	Equate derivative to zero and solve for x	DM1	
	Obtain $x = \frac{1}{a}$, $y = \frac{1}{ae}$	A1	ISW Or exact equivalent.
		4	
6(b)	Use integration by parts to obtain $pxe^{-ax} + q \int e^{-ax} dx$	*M1	Condone sign error in parts formula and omission of dx . M0 if a value is used for a (not equivalent work).
	Obtain $-\frac{1}{a}xe^{-ax} + \frac{1}{a} \int e^{-ax} dx$	A1	OE
	Complete integration to obtain $-\frac{1}{a}xe^{-ax} - \frac{1}{a^2}e^{-ax}$	A1	OE
	Correct use of limits 0 and $\frac{2}{a}$ in an expression of the form $rx e^{-ax} + se^{-ax}$	DM1	$\left(\frac{-2}{a^2}e^{-2} - \frac{1}{a^2}e^{-2} + 0 + \frac{1}{a^2}\right)$
	Obtain $\frac{1}{a^2}(1 - 3e^{-2})$	A1	ISW Or simplified 2-term equivalent, e.g. $\frac{e^2 - 3}{a^2 e^2}$.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	Factorise LHS using difference of 2 squares	*M1	$((\cos^2 \theta - \sin^2 \theta)(\cos^2 \theta + \sin^2 \theta))$
	Simplify	DM1	$\cos^2 \theta + \sin^2 \theta = 1$ must be seen or implied, e.g. $(\cos^2 \theta - \sin^2 \theta)(\cos^2 \theta + \sin^2 \theta) = (\cos^2 \theta - \sin^2 \theta)$.
	Obtain $\cos^4 \theta - \sin^4 \theta \equiv \cos 2\theta$ from correct working	A1	AG
	Alternative Method for Question 7(a)		
	Use of correct rearrangements of double angle formulae	(*M1)	E.g. $\left(\frac{1 + \cos 2\theta}{2}\right)^2 - \left(\frac{1 - \cos 2\theta}{2}\right)^2$ Only condone $\left(\frac{1 + \cos 2\theta}{2}\right)^2 - \left(\frac{\cos 2\theta - 1}{2}\right)^2$ if correct expression for $\sin^2 \theta$ seen.
	Expand and simplify	(DM1)	Collect like terms. Condone recovery from missing brackets.
	Obtain $\cos^4 \theta - \sin^4 \theta \equiv \cos 2\theta$ from correct working	(A1)	AG
	Alternative Method 2 for Question 7(a)		
	Correct use of Pythagoras	(*M1)	E.g. $(1 - \sin^2 \theta)^2 - \sin^4 \theta$ or $\cos^2 \theta(1 - \sin^2 \theta) - \sin^2 \theta(1 - \cos^2 \theta)$
	Expand and simplify	(DM1)	Condone recovery from missing brackets.
Obtain $\cos^4 \theta - \sin^4 \theta \equiv \cos 2\theta$ from correct working	(A1)	AG	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
7(b)	Use part (a) and correct double angle formula to obtain expression involving $\int \sin^2 2\theta d\theta$ or $\int \cos^2 2\theta d\theta$	M1	$\int \cos^4 \theta - \sin^4 \theta + 4\sin^2 \theta \cos^2 \theta d\theta = \int \cos 2\theta + \sin^2 2\theta d\theta$ Allow BOD for $2\sin^2 2\theta$ if $\sin 2\theta = 2\sin \theta \cos \theta$ seen.
	$\int \cos 2\theta d\theta = \frac{1}{2} \sin 2\theta$	B1	Seen or implied.
	Use of correct double angle formula on second part of the integral to obtain a form that can be integrated directly	M1	e.g. $\int \sin^2 2\theta d\theta = \int \frac{1 - \cos 4\theta}{2} d\theta$
	Obtain $\frac{1}{2}\theta - \frac{1}{8}\sin 4\theta$	A1	Condone a mixture of x and θ .
	Correct use of limits $\pm \frac{\pi}{8}$ in an expression of the form $p\theta + q\sin 2\theta + r\sin 4\theta$ and evaluate the trig	M1	$\left(2\left(\frac{1}{2} \times \frac{1}{\sqrt{2}} + \frac{\pi}{16} - \frac{1}{8}\right)\right)$
	Obtain $\frac{1}{2}\sqrt{2} + \frac{1}{8}\pi - \frac{1}{4}$	A1	ISW Or exact equivalent from exact working.
			6


PUBLISHED

Question	Answer	Marks	Guidance
8(a)	Correct direction vector seen or implied ($\overline{BC} = 3\mathbf{i} + 3\mathbf{j} - 3\mathbf{k}$)	B1	Condone $\overline{BC} = -3\mathbf{i} - 3\mathbf{j} + 3\mathbf{k}$.
	Use a correct method to form a vector equation	M1	Allow for the RHS with no LHS.
	Obtain $\mathbf{r} = 5\mathbf{i} + 2\mathbf{j} + \lambda(\mathbf{i} + \mathbf{j} - \mathbf{k})$	A1	ISW Must have $\mathbf{r} = \dots$ or $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \dots$, not $l_1 = \dots$ Or, equivalent vector form, e.g. $\mathbf{r} = 8\mathbf{i} + 5\mathbf{j} - 3\mathbf{k} + \alpha(\mathbf{i} + \mathbf{j} - \mathbf{k})$ or $\mathbf{r} = 5\mathbf{i} + 2\mathbf{j} + \lambda(3\mathbf{i} + 3\mathbf{j} - 3\mathbf{k})$. Condone a column vector with $\mathbf{i}, \mathbf{j}, \mathbf{k}$.
		3	
8(b)	Use components to form two relevant equations in 2 unknowns For their l_1 B0 if they use the same unknown for both lines.	B1FT	Two components of $\begin{pmatrix} 5 + \lambda \\ 2 + \lambda \\ -\lambda \end{pmatrix} = \begin{pmatrix} -2 + 3\mu \\ 1 + \mu \\ 4 - 2\mu \end{pmatrix}$ seen or implied.
	Solve 2 relevant equations in 2 unknowns for λ or μ	M1	For <i>their</i> l_1 .
	Obtain $\lambda = 2$ or $\mu = 3$	A1	Or equivalent e.g. using \overline{BC} as direction vector gives $\lambda = \frac{2}{3}$.
	Obtain $(7, 4, -2)$ No need to check the third equation – the question implies that the lines intersect.	A1	Accept position vector. Condone a column vector with $\mathbf{i}, \mathbf{j}, \mathbf{k}$. SC: B1 M1 A1 A1 if one component of their line is incorrect but they do not use that component.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
8(c)	State $AB = \sqrt{7^2 + 1^2 + 4^2} (= \sqrt{66})$	B1	Or $(AB)^2 = 66$ Condone a sign error in \overline{AB} .
	State \overline{BD} in component form	B1	$\begin{pmatrix} -7 + 3r \\ -1 + r \\ 4 - 2r \end{pmatrix}$ or equivalent.
	$AB = BD \Rightarrow (3r - 7)^2 + (r - 1)^2 + (-2r + 4)^2 = 66$ $(14r^2 - 60r = 0)$	M1	Or equivalent equation in one unknown for their AB and <i>their</i> $\overline{BD} \neq \overline{OD}$. If you never see a correct form and they go direct to $9r^2 + 49 + r^2 + 1 \dots$ then M0.
	$\Rightarrow r = \frac{30}{7}$	A1	Correct only. Ignore $r = 0$ if seen.
	$\overline{OD} = \frac{76}{7}\mathbf{i} + \frac{37}{7}\mathbf{j} - \frac{32}{7}\mathbf{k}$	A1	Must be a vector. Condone if also have $\overline{OD} = \overline{OA}$.
		5	

Question	Answer	Marks	Guidance
9(a)	State $z\omega = (-3 + 3\sqrt{3}) + (3 + 3\sqrt{3})\mathbf{i}$	B1	Or exact equivalent with real and imaginary parts collected. Need brackets around the coefficient of \mathbf{i} . Allow for $a =$, $b =$ stated correctly.
		1	

Question	Answer	Marks	Guidance
9(b)	Obtain $ z = \sqrt{2}$	B1	
	Obtain $\arg z = -\frac{\pi}{4}$ final answer	B1	
	Obtain $ \omega = 6$	B1	
	Obtain $\arg \omega = \frac{2\pi}{3}$ final answer	B1	
		4	
9(c)			<p>Note: The question does not require the diagram. If they use $\frac{5\pi}{12}$ they need to demonstrate where it comes from. Complex number equivalent to AB is $3\sqrt{3} + 3i$.</p>
	Show $ OA = AB = 6$, hence isosceles	B1	One mark for ‘isosceles’ and one mark for ‘right angle’.
	$\angle AOB = \arg \omega - \arg z = -\arg z = \frac{\pi}{4}$ hence third angle is a right angle	B1	There will be alternatives e.g. use of Pythagoras (ratio of lengths is $1:1:\sqrt{2}$), expressing each number in “vector” form and using scalar product or explaining the effect of multiplying by $1 - i$.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
9(d)	$\arg z\omega = \arg z + \arg \omega \left(= \frac{2\pi}{3} - \frac{\pi}{4} = \frac{5\pi}{12} \right)$	M1	For showing correct use of their angles from part (b). Must demonstrate where $\frac{5\pi}{12}$ comes from.
	$\arg z\omega = \tan^{-1} \frac{3+3\sqrt{3}}{-3+3\sqrt{3}}$	M1	Correct method for <i>their</i> $z\omega$ from part (a). Must link to point <i>B</i> on diagram or to $\arg z\omega$. Need to see $\tan^{-1} \frac{3+3\sqrt{3}}{-3+3\sqrt{3}}$ or $\tan \theta = \frac{3+3\sqrt{3}}{-3+3\sqrt{3}}$ and not just $\tan^{-1} \frac{1+\sqrt{3}}{-1+\sqrt{3}}$.
	$\Rightarrow \tan\left(\frac{5}{12}\pi\right) = \frac{\sqrt{3}+1}{\sqrt{3}-1}$	A1	Obtain given answer from full and correct working.
		3	

Question	Answer	Marks	Guidance
10(a)	Use of correct chain rule (and correct quotient rule) and $\cos^{-3} \theta$	M1	Obtain $k \times (\cos \theta)^{-4} \times \sin \theta$ or equivalent.
	$\frac{dy}{d\theta} = -3 \times -\sin \theta (\cos \theta)^{-4} = 3 \sin \theta \sec^4 \theta$ Must be expressed in the given form	A1	Obtain given answer from full and correct working (signs must be shown), but condone $\frac{d}{d\theta} (\sec^3 \theta) = \dots$ and $y'(\theta)$.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
10(b)	Separate variables: $\int \frac{\sin \theta}{\cos^4 \theta} d\theta = \int \frac{(x+3)}{(x^2+9)} dx$	B1	Or $\int \frac{3 \sin \theta}{\cos^4 \theta} d\theta = \int \frac{3x+9}{x^2+9} dx$. Condone missing integral signs or missing dx or dθ, but not both.
	Obtain $p \sec^3 \theta (+A)$	B1	Correct form, p any constant but not 0.
	Use $\int \frac{3x+9}{x^2+9} dx = \int \left(\frac{3x}{x^2+9} + \frac{9}{x^2+9} \right) dx$ and obtain $q \ln(x^2+9)$ or $r \tan^{-1} \frac{x}{3} (+C)$.	*M1	Might have one third of both sides. Alt: substitute $x = 3 \tan \phi$ to obtain $q \int 1 + \tan \phi d\phi$; condone if have θ in place of ϕ in this method.
	Obtain $q \ln(x^2+9)$ and $r \tan^{-1} \frac{x}{3} (+C)$	DM1	Obtain $q(\phi \mp \ln(\cos \phi))$ OE.
	Obtain $\sec^3 \theta = \frac{3}{2} \ln(x^2+9) + 3 \tan^{-1} \frac{x}{3} (+C)$ or equivalent	A1	Or might see a third of both sides. Must have 2 different variables.
	Use $\theta = \frac{1}{3}\pi$, $x = 3$ in an equation including $p \sec^3 \theta$, $q \ln(x^2+9)$ and $r \tan^{-1} \frac{x}{3}$ to evaluate the constant of integration	M1	Or as limits in a definite integral. Limits for ϕ are 0 and $\frac{1}{4}\pi$.
	Obtain constant = $8 - \frac{3}{2} \ln 18 - \frac{3}{4}\pi$	A1	OE, e.g. 1.308... to at least 3sf.
	Obtain $\cos \theta = 0.601$	A1	Accept AWRT 0.601.
			8